

SHORT REPORT

Open Access

A non-radioactive method for small RNA detection by northern blotting

Qi Huang^{1,2}, Zhinang Mao¹, Shaoqing Li^{1,2}, Jun Hu^{1,2,3*} and Yingguo Zhu^{1,2*}

Abstract

Background: Small non-coding RNAs are essential regulators of gene expression at the transcriptional and posttranscriptional levels. High-throughput sequencing has revealed thousands of predicted small RNAs; however, only a few of these have been well characterized. Northern blotting is the most convincing method for small RNA validation.

Findings: In this study, we improved the Northern blot method by using biotin-labeled probes. miRNAs and siRNAs derived from both *Arabidopsis thaliana* and *Oryza sativa* were investigated. The results suggest that this improved method is sensitive and efficient, with approximately 5 µg of total RNA being sufficient for detection. Furthermore, long-term storage of probes labeled in this manner is more convenient, less contaminative and degradative compared with traditional probes.

Conclusions: This protocol is an alternative strategy for small RNA detection and represents an efficient means of researching small RNAs.

Keywords: Small RNA; Biotin-labeled probe; Non-radioactive; Northern blot; Epigenetic

Findings

In eukaryotic cells, small non-coding RNAs have been demonstrated to play fundamental roles in gene expression modification during development (Zamore & Haley, 2005). Small non-coding RNAs are 20-30 nt long and function as sequence-specific negative regulators of gene expression at the transcriptional and/or posttranscriptional levels (He & Hannon, 2004; Voinnet, 2009). The three major types of small non-coding RNAs—miRNAs, siRNAs, and piRNAs—are distinguished by their different modes of biogenesis (Chen, 2009; Huang et al, 2011). In plants, miRNAs and siRNAs are the two major classes of endogenous small RNAs and are important for development, genome stability, gene expression and stress responses (Chen, 2012; Wei et al, 2012). Although the mechanisms and pathways by which miRNAs and siRNAs regulate their target genes are largely obscure, the initial findings have demonstrated that miRNAs and siRNAs have a broad impact on epigenetics.

Since the discovery of miRNAs in *C. elegans* two decades ago, remarkable advances in the characterization of this gene family have been achieved (He & Hannon, 2004). Recently, high-throughput sequencing technology has facilitated the exploration of small non-coding RNAs (Creighton et al, 2009; Studholme, 2012). To date, thousands of small non-coding RNAs and their target mRNAs or genes have been computationally predicted; however, few of them have been experimentally confirmed or properly characterized. To gain further insight, several methods have been developed to investigate the expression of target small non-coding RNAs, such as Northern blotting, quantitative reverse-transcription PCR (RT-PCR), and in situ hybridization.

rRNAs and tRNAs account for over 90% of the total RNA in a eukaryotic cell, whereas small non-coding RNAs account for only approximately 2%, and detection of these RNAs by some methods is difficult (Zhuang et al, 2012). Because the low abundance of small RNAs can be problematic for detection via northern blotting, some researchers have enriched small RNAs using LiCl or other methods (Song et al, 2012). Although isotope labeling is often inconvenient, hazardous and restricted by many institutions, this classic method is still the most popular

* Correspondence: junhu@whu.edu.cn; zhu yg@public.wh.hb.cn

¹State Key Laboratory of Hybrid Rice, College of Life Sciences, Wuhan University, Wuhan 430072, China

²Engineering Research Center for Plant Biotechnology and Germplasm, Utilization, Ministry of Education, Wuhan University, Wuhan 430072, China
Full list of author information is available at the end of the article

method for investigating the expression of small RNAs. Several improvements have been made to traditional Northern blotting protocols, and non-isotopic-labeling methods using digoxigenin (DIG) represent a safe alternative for the detection of small RNAs (Ramkissoon et al, 2006). In recent years, researchers have been continuously improving the method for northern blotting with digoxigenin (DIG)-labeled modified probes (Kim et al, 2010). Researchers have also modified probes with locked nucleic acid (LNA) to improve sensitivity (Gao & Peng, 2011; Lopez-Gomollon, 2011).

Biotin is another molecule that can increase the sensitivity of small RNA detection. In this study, we designed biotin-labeled oligonucleotide probes to investigate the expression of small RNAs. When we applied this improved method to *Arabidopsis thaliana* and *Oryza sativa* for validation, the results suggested that this method was sensitive and efficient, as it was capable of detecting as little as 1 to 5 μg of total RNA. Furthermore, storage of biotin-labeled probes is convenient, and the probes are stable; therefore, northern blots can be carried out with ease. Taken together, these data suggest that biotin-labeled probes can be widely used to investigate small RNAs and even mRNAs in the future.

Preparation of total RNA and small RNA for Northern blotting

In a eukaryotic cell, the abundance of rRNA and tRNA impairs the detection of small RNAs. To test the method of RNA preparation in this study, total RNA and small RNAs were extracted with two different reagents. To detect the integrity of the RNA, we loaded the RNA onto 15% PAGE gels under denaturing conditions (7 M urea) or onto 1.2% agarose gels with EtBr for staining. The results showed that the RNAs were suitable for further analysis, such as northern blotting (Figure 1a,b). Next,

the amounts of RNA were quantified using a NanoDrop (ThermoScientific). The data suggested that the small RNAs, including 5S rRNA, tRNA and small non-coding RNAs, accounted for approximately 25% of the total RNA, and three-quarters of the unwanted RNA was removed effectively (Table 1). Hence, this RNA isolation method is suitable for extracting small fragment RNAs and for removing the abundant RNAs. Northern blots were further performed to detect the expression of miRNAs. miRNA5078 was specifically investigated using a miRNA5078-biotin-labeled probe. The data established that miRNA5078 is present in leaves (Figure 1c). This result also indicated that 5 μg of total RNA or 1 μg of small RNA was sufficient for detection.

Northern blots of miRNA and siRNA of *Arabidopsis*

Because northern blots performed using biotin-labeled probes were sensitive and efficient, we used this method to confirm the expression of classic miRNAs and siRNAs in *Arabidopsis*. It is well known that the abundance of small non-coding RNAs is less than 2%. To address this problem, researchers usually increase the amount of input RNA to 20–60 μg depending on the expression level of the target miRNA. In this study, the input RNAs (ranging from 1 to 5 μg) were loaded onto 15% denatured PAGE gels before northern blotting to test the sensitivity of this method. We examined certain miRNAs (miR156, miR171a, miR390b, and miR168a) because they had previously been shown to be expressed in leaves (Yang et al, 2006; Zhan et al, 2012). As shown in Figure 2a, these miRNAs were easily detected after a 1 to 5 minute exposure time, which was determined based on the typical amount of miRNA *in vivo*. Four gene families have been identified in *Arabidopsis* that produce a number of trans-acting siRNAs (ta-siRNAs) (Allen et al, 2005). ta-siRNAs, such as tasiR-ARF, also play essential




Figure 1 Integrity of extracted RNAs. (a) Total RNA (left) and small RNAs (right) were loaded on 15% denaturing PAGE gels and stained with 0.5 $\mu\text{g}/\text{ml}$ EtBr in buffer. **(b)** Total RNA (left) and small RNAs (right) were loaded on 1.2% agarose gels and stained with 0.5 $\mu\text{g}/\text{ml}$ EtBr in buffer. **(c)** Northern blot of miRNA5078 with variable amounts of total RNA (left) and small RNAs (right). The exposure times are indicated.

Table 1 Valuation of extracted RNAs

	Total RNAs (Leaves)	Small RNAs (Leaves)
Concentration (µg/ml)	2105.0 ± 24.5	538.5 ± 16.4
OD260:280	2.01	1.98

Extracted RNAs were quantified with NanoDrop 2000c Spectrophotometer. Values are means ± S.D.

roles in plant development (Williams *et al*, 2005). Subsequently, we also evaluated the accumulation of ta-siRNAs, such as ta-siRNA2141, ta-siRNA752, ta-siRNA255, and ta-siRNA850. The results demonstrate that this method is also efficient for studying the expression of ta-siRNAs (Figure 2b). Therefore, we confirmed that both miRNAs ta-siRNAs can be detected by this method with small quantities of total RNA.

Northern blotting of miRNAs and ta-siRNAs from rice

Rice is an important staple crop worldwide, and it is also an essential model crop for research. To ensure that this method can be applied to rice, we also validated established rice miRNAs and ta-siRNAs. Based on previous studies and the Plant MicroRNA Database (<http://bioinformatics.cau.edu.cn/PMRD/>), miR156, miR171a, miR390b, and miR168a were selected for investigation. As shown in Figure 3a, most miRNAs were easily detected in 5 µg of total RNA with 1 minute of exposure time. Furthermore, the results suggested that detection of ta-siRNA was also successful (Figure 3b). These results demonstrate that this method is an efficient means of assessing the presence of small RNAs in small amounts of total RNA. Indeed, 1 µg of total RNA was sufficient for the detection of some highly expressed small RNAs. For small RNAs with lower expression levels, 5 µg of total RNA was sometimes necessary for detection. Consequently, 1 µg of purified small RNA is sufficient for northern blot applications with this strategy, and this amount can easily be obtained by researchers without any special or expensive equipment.

Protocol for northern blotting with biotin-labeled probes

The methodology described in this study is illustrated in Figure 4. First, the integrity of the RNAs is determined, and the total amount is quantified. Both total RNA and small fragment RNAs can be used for detection, and the amount of the latter needed for detection was only approximately one-quarter of the total RNA. If the precursors of miRNAs or ta-siRNAs were not needed for research, small fragment RNAs were preferred as the loading sample. To avoid degradation of the RNA, the glass plates used for RNA gels should first be wiped with ethanol. It is recommended that the 15% denaturing PAGE gel containing 7 M urea and TBE running buffer be prepared with DEPC treated water. The biotin can be modified on either the 5' or 3' terminus, as both methods are available. Probes can be stored as 100 µM stocks at 4°C for 6 months or at -20°C for long-term storage. The complete protocol requires only two days, including overnight hybridization on the first day.

Discussion

The massive increase in the number of small RNAs being uncovered in numerous organisms increases the need for improved methods of small RNA detection. The overwhelming majority of studies on small RNAs suggest that northern blotting is the most popular method for studying the expression of target small RNAs. However, isotope contamination is a serious problem for the environment and researcher safety, and most researchers are eager to find a safer alternative. Digoxigenin, which is widely used for Southern blots, northern blots, and in situ hybridization, may represent a safe alternative label. We also acknowledge the discovery of biotin, which makes such research easier and more convenient.

In addition to isotope safety, sensitivity is another problem associated with northern blotting. Several improved




Figure 2 Detection of miRNAs and ta-siRNAs from *Arabidopsis*. (a) Northern blots for miRNAs using biotin-labeled probes and variable amounts of total RNA from *Arabidopsis* leaves. The exposure time is indicated. (b) Northern blots for ta-siRNAs using biotin-labeled probes and variant total RNA from *Arabidopsis* leaves. The exposure time is indicated.




Figure 3 Detection of miRNAs and ta-siRNAs from rice. **(a)** Northern blots for miRNAs using biotin-labeled probes and variant total RNA from rice leaves. The exposure time is indicated. **(b)** Northern blots for ta-siRNAs using biotin-labeled probes and variant total RNA from rice leaves. The exposure time is indicated.

technologies have been reported, including the use of chemical cross-linking instead of UV, the use of an LNA primer as a probe (Valoczi et al, 2004), LED (LNA, EDC, DIG) modification (Kim et al, 2010) and purification and loading of only the small RNA fraction (Song et al, 2012). Improving the sensitivity of these methods is important, as the amount of miRNA or siRNA is often very low. Biotin is becoming more widely used in biotechnological applications. The affinity of biotin for streptavidin is one of the strongest and most stable non-covalent interactions. Horseradish Peroxidase (HRP) is an enzyme that utilizes organic peroxide compounds as electron donors. The biotin-streptavidin-HRP conjugation method is ideal for investigators working in the fields of immunology, cancer, and neuroscience as well as other areas.

In this study, we significantly improved the northern blotting procedure for detecting small RNAs with the use of a biotin-labeled probe. Biotin-labeled probes have the benefits of being safer and easier to handle and store compared with traditional probes. A biotin-labeled probe can be stored for at least 6 months at 4°C, whereas a ³²P-labeled probe can be stored for no more than 1 month. Furthermore, biotin-labeled probes are synthesized once and can be re-used more than 100 times, whereas isotope-labeled probes must be re-labeled with every use. Probe sensitivity is extremely important, and the sensitivity of biotin-labeled probes is acceptable based on our studies. Typically, 20–40 µg total RNA is used for northern blots, and even for rice, some researchers still use 30 µg of small fragment RNAs or total RNA for detection by isotope-labeled probes (Wang et al, 2004; Zhong et al, 2013). When using biotin-labeled probes, the amount of input RNA can be decreased to 2–5 µg based on our research. If the amount of a particular target small RNA is very low, we recommend increasing the amount of total RNA to 20 µg and the amount of small

fragment RNAs to 5 µg. Biotin probes can also save time, as only two days are needed to complete the procedure, and there is no need to spend time labeling or purifying isotopes. The results from this study using both *Arabidopsis* and rice suggest that this method works well and may be suitable for use in other species.

Conclusions

To date, high-throughput sequencing has revealed thousands of predicted small RNAs; however, only a few of them have been characterized. Most researchers use isotope-labeled probes for small RNA detection and validation. Although biotin has been used in many fields, Northern blots using biotin-labeled probes have not been reported. In this study, we improved Northern blotting with the use of biotin-labeled probes. Northern blots using RNA derived from both *Arabidopsis* and *Oryza sativa* were performed, and the results suggest that this improved method is both sensitive and efficient. Furthermore, biotin-labeled probes are more stable and safer for researchers and the environment than traditional probes. This protocol represents an important contribution to the small RNA field.

Materials and methods

Probe Preparation

Some of the miRNAs and ta-siRNAs selected for validation are listed in the small RNA database (<http://bioinformatics.cau.edu.cn/PMRD/>), and some have been reported in previous studies (Allen et al, 2005; Xie et al, 2006; Zhan et al, 2012). Probes modified with biotin on the 5' or 3' terminus were synthesized and purified via HPLC by the GenScript Company (Nanjing, China). These probes were designed to be completely complementary to the miRNA or ta-siRNA nucleotide sequence and are listed in Additional file 1: Table S1.




Figure 4 Schematic diagram of Northern blotting with biotin-labeled probes.

RNA extraction

RNA was extracted from the leaves of *Arabidopsis* and *Oryza sativa*. Total RNA was extracted from leaves using TRIzol reagent (Invitrogen, USA) following a previously described protocol (Hu et al, 2012). Small RNAs were extracted from leaves using RNAiso for Small RNA (TaKaRa, Dalian, China). Approximately 50 mg of leaf tissue was ground into a powder under liquid nitrogen and incubated with 1 ml of RNAiso reagent. The tubes were shaken vigorously by hand for 30 seconds and incubated for 5 minutes prior to centrifugation at 12,000 g for 15 minutes at 4°C. The aqueous (upper) phase of the sample was transferred to a new RNase-free tube by carefully angling the tube at 45°. The aqueous phase was extracted with chloroform by vortexing for 2 minutes and centrifuging the mixture at 12,000 g for 5 minutes

at 4°C. Small RNAs were precipitated with an equal volume of isopropanol at -20°C for at least 1 hour, followed by centrifugation at 12,000 g for 15 minutes at 4°C. The RNA pellets were washed with 200 µl of 75% ethanol and resuspended in RNase-free water. Finally, total RNA and small RNAs were quantified using a NanoDrop 2000c Spectrophotometer (Thermo Scientific, USA) and visually assessed on a 15% denaturing PAGE gel containing 7 M urea or on a 1.2% agarose gel.

RNA transfer and UV cross-linking

The 15% denatured polyacrylamide gels containing 7 M urea were 15 cm wide, 20 cm long, and 1.5 mm thick. After polymerization at room temperature for 30 minutes, the gels were pre-run with 1× TBE at 40 mA (600 V) for 30 minutes. The RNAs or oligonucleotide probes were mixed

with 2× loading buffer (5 mM EDTA, 0.1% bromophenol blue, 0.1% xylene cyanol, and 95% formamide); the RNA samples were heated at 70°C for 5 minutes. The PAGE gels were run at 40 mA (600 V) for approximately 2 hours until the bromophenol blue reached approximately 1 cm above the bottom of the gel. The gels were then stained with 0.5 µg/ml EtBr in DEPC-treated TBE buffer for 10 minutes. The RNAs were then transferred onto Hybond-N + positively charged nylon membranes (GE healthcare, USA) using a Bio-Rad transblot apparatus at 200 mA (9 - 10 V) for 3 hours (Bio-Rad, USA). Next, the membranes were further crosslinked at 1200 µjoules for 20 minutes and dried at 50°C for 30 minutes to improve sensitivity. Membranes prepared in this way can be stored at 4°C for several months.

Hybridization

Before hybridization, the membranes were pre-hybridized for at least 30 minutes at 40°C in pre-hybridization buffer (7% SDS, 200 mM Na₂HPO₄ (pH 7.0), 5 µg/ml salmon sperm DNA (SSDNA)). A 3-hour incubation period is recommended for decreasing background noise. Next, the pre-hybridization buffer was removed, and hybridization buffer containing 50 pmol/ml labeled probes was added. The biotin-labeled probe did not require denaturation. The membranes were hybridized for 12 to 16 hours at 40°C with gentle shaking and subsequently rinsed with washing buffer (1× SSC, 0.1% SDS) 3 times for approximately 15 minutes total at room temperature.

ECL lighting

The biotin-labeled probes were detected using a Chemiluminescent Nucleic Acid Detection Module Kit (Thermo Scientific, USA). After rinsing, the membranes were blocked again with blocking buffer for 15 minutes with gentle shaking at room temperature followed by incubation for an additional 15 minutes with hybridization buffer containing stabilized streptavidin-HRP conjugate. After washing 3 times, the membranes were equilibrated in substrate equilibration buffer for 5 minutes. Finally, the membranes were placed in a clean container, covered completely with working solution, and incubated for 5 minutes in the dark. A working solution was prepared with luminol/enhancer solution and stable peroxide solution. The membranes were placed in a cassette with X-ray films and exposed for different periods of time (depending on the desired signal intensity).

Additional file

Additional file 1: Table S1. Probes used in this study.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

QH performed the experiments and analyzed the data, ZM performed the experiments and analyzed the data, SL performed the experiments and analyzed the data, JH conceived the study, performed the experiments, analyzed the data and wrote the manuscript, YZ conceived the study. All authors read and approved the final manuscript.

Acknowledgments

We greatly thank Dr. Qijia Wu and Dr. Xiaorong Zhang (Wuhan University) and Professor Yi Zhang (ABLife Company, Wuhan) for technical assistance and critical comments. This study was supported by funds from the National High-Tech Research and Development Project (20011AA101101), the National Key Project for Research on Transgenic Plants (2011ZX08010), and the National Natural Science Foundation of China (31371698).

Author details

¹State Key Laboratory of Hybrid Rice, College of Life Sciences, Wuhan University, Wuhan 430072, China. ²Engineering Research Center for Plant Biotechnology and Germplasm, Utilization, Ministry of Education, Wuhan University, Wuhan 430072, China. ³Suzhou institute of Wuhan University, Wuhan 215000, China.

Received: 1 August 2014 Accepted: 20 September 2014

Published online: 01 October 2014

References

- Allen E, Xie Z, Gustafson AM, Carrington JC (2005) microRNA-directed phasing during trans-acting siRNA biogenesis in plants. *Cell* 121:207–221
- Chen X (2009) Small RNAs and their roles in plant development. *Annu Rev Cell Dev Biol* 25:21–44
- Chen X (2012) Small RNAs in development - insights from plants. *Curr Opin Genet Dev* 22:361–367
- Creighton CJ, Reid JG, Gunaratne PH (2009) Expression profiling of microRNAs by deep sequencing. *Brief Bioinform* 10:490–497
- Gao Z, Peng Y (2011) A highly sensitive and specific biosensor for ligation- and PCR-free detection of MicroRNAs. *Biosens Bioelectron* 26(9):3768–3773
- He L, Hannon GJ (2004) MicroRNAs: small RNAs with a big role in gene regulation. *Nat Rev Genet* 5:522–531
- Hu J, Wang K, Huang W, Liu G, Gao Y, Wang J, Huang Q, Ji Y, Qin X, Wan L, Zhu R, Li S, Yang D, Zhu Y (2012) The rice pentatricopeptide repeat protein RF5 restores fertility in Hong-Lian cytoplasmic male-sterile lines via a complex with the glycine-rich protein GRP162. *Plant Cell* 24:109–122
- Huang H, Gao Q, Peng X, Choi SY, Sarma K, Ren H, Morris AJ, Frohman MA (2011) piRNA-associated germline nuage formation and spermatogenesis require MitoPLD profusogenic mitochondrial-surface lipid signaling. *Dev Cell* 20:376–387
- Kim SW, Li Z, Moore PS, Monaghan AP, Chang Y, Nichols M, John B (2010) A sensitive non-radioactive northern blot method to detect small RNAs. *Nucleic Acids Res* 38:e98
- Lopez-Gomollon S (2011) Detecting sRNAs by Northern blotting. *Methods Mol Biol* 732:25–38
- Ramkisson SH, Mainwaring LA, Sloand EM, Young NS, Kajigaya S (2006) Nonisotopic detection of microRNA using digoxigenin labeled RNA probes. *Mol Cell Probes* 20:1–4
- Song X, Wang D, Ma L, Chen Z, Li P, Cui X, Liu C, Cao S, Chu C, Tao Y, Cao X (2012) Rice RNA-dependent RNA polymerase 6 acts in small RNA biogenesis and spikelet development. *Plant J* 71:378–389
- Studholme DJ (2012) Deep sequencing of small RNAs in plants: applied bioinformatics. *Brief Funct Genomics* 11:71–85
- Valoczi A, Hornyik C, Varga N, Burgyan J, Kauppinen S, Havelda Z (2004) Sensitive and specific detection of microRNAs by northern blot analysis using LNA-modified oligonucleotide probes. *Nucleic Acids Res* 32:e175
- Voynet O (2009) Origin, biogenesis, and activity of plant microRNAs. *Cell* 136:669–687
- Wang JF, Zhou H, Chen YQ, Luo QJ, Qu LH (2004) Identification of 20 microRNAs from *Oryza sativa*. *Nucleic Acids Res* 32:1688–1695
- Wei W, Ba Z, Gao M, Wu Y, Ma Y, Amiard S, White CI, Rendtlew Danielsen JM, Yang YG, Qi Y (2012) A role for small RNAs in DNA double-strand break repair. *Cell* 149:101–112

- Williams L, Carles CC, Osmont KS, Fletcher JC (2005) A database analysis method identifies an endogenous trans-acting short-interfering RNA that targets the Arabidopsis ARF2, ARF3, and ARF4 genes. *Proc Natl Acad Sci U S A* 102:9703–9708
- Xie K, Wu C, Xiong L (2006) Genomic organization, differential expression, and interaction of SQUAMOSA promoter-binding-like transcription factors and microRNA156 in rice. *Plant Physiol* 142:280–293
- Yang L, Liu Z, Lu F, Dong A, Huang H (2006) SERRATE is a novel nuclear regulator in primary microRNA processing in Arabidopsis. *Plant J* 47:841–850
- Zamore PD, Haley B (2005) Ribo-gnome: the big world of small RNAs. *Science* 309:1519–1524
- Zhan X, Wang B, Li H, Liu R, Kalia RK, Zhu JK, Chinnusamy V (2012) Arabidopsis proline-rich protein important for development and abiotic stress tolerance is involved in microRNA biogenesis. *Proc Natl Acad Sci U S A* 109:18198–18203
- Zhong SH, Liu JZ, Jin H, Lin L, Li Q, Chen Y, Yuan YX, Wang ZY, Huang H, Qi YJ, Chen XY, Vaucheret H, Chory J, Li J, He ZH (2013) Warm temperatures induce transgenerational epigenetic release of RNA silencing by inhibiting siRNA biogenesis in Arabidopsis. *Proc Natl Acad Sci U S A* 110:9171–9176
- Zhuang F, Fuchs RT, Robb GB (2012) Small RNA expression profiling by high-throughput sequencing: implications of enzymatic manipulation. *Nucleic Acids* 2012:360358

doi:10.1186/s12284-014-0026-1

Cite this article as: Huang *et al.*: A non-radioactive method for small RNA detection by northern blotting. *Rice* 2014 **7**:26.

Submit your manuscript to a SpringerOpen[®] journal and benefit from:

- ▶ Convenient online submission
- ▶ Rigorous peer review
- ▶ Immediate publication on acceptance
- ▶ Open access: articles freely available online
- ▶ High visibility within the field
- ▶ Retaining the copyright to your article

Submit your next manuscript at ▶ springeropen.com
